



Inter-Individual Variability of Stone Marten Behavioral Responses to a Highway

Fernando Ascensão^{1,2*}, Clara Grilo³, Scott LaPoint^{4,5}, Jeff Tracey⁶, Anthony P. Clevenger², Margarida Santos-Reis¹

1 Centro de Biologia Ambiental, Faculdade de Ciências da Universidade de Lisboa, Lisboa, Portugal, **2** Western Transportation Institute, Montana State University, Bozeman, Montana, United States of America, **3** Departamento de Biologia & CESAM, Universidade de Aveiro, Aveiro, Portugal, **4** Max-Planck-Institute for Ornithology, Radolfzell, Germany, **5** Department of Biology, University of Konstanz, Konstanz, Germany, **6** US Geological Survey, Western Ecological Research Center, San Diego, California, United States of America

Abstract

Efforts to reduce the negative impacts of roads on wildlife may be hindered if individuals within the population vary widely in their responses to roads and mitigation strategies ignore this variability. This knowledge is particularly important for medium-sized carnivores as they are vulnerable to road mortality, while also known to use available road passages (e.g., drainage culverts) for safely crossing highways. Our goal in this study was to assess whether this apparently contradictory pattern of high road-kill numbers associated with a regular use of road passages is attributable to the variation in behavioral responses toward the highway between individuals. We investigated the responses of seven radio-tracked stone martens (*Martes foina*) to a highway by measuring their utilization distribution, response turning angles and highway crossing patterns. We compared the observed responses to simulated movement parameterized by the observed space use and movement characteristics of each individual, but naïve to the presence of the highway. Our results suggested that martens demonstrate a diversity of responses to the highway, including attraction, indifference, or avoidance. Martens also varied in their highway crossing patterns, with some crossing repeatedly at the same location (often coincident with highway passages). We suspect that the response variability derives from the individual's familiarity of the landscape, including their awareness of highway passage locations. Because of these variable yet potentially attributable responses, we support the use of exclusionary fencing to guide transient (e.g., dispersers) individuals to existing passages to reduce the road-kill risk.

Citation: Ascensão F, Grilo C, LaPoint S, Tracey J, Clevenger AP, et al. (2014) Inter-Individual Variability of Stone Marten Behavioral Responses to a Highway. PLoS ONE 9(7): e103544. doi:10.1371/journal.pone.0103544

Editor: Maharaj K. Pandit, University of Delhi, India

Received: March 7, 2014; **Accepted:** July 4, 2014; **Published:** July 29, 2014

Copyright: © 2014 Ascensão et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. not applicable because we reanalyzed data previously published. Simulation code will be as a Supporting Information file.

Funding: FA and CG were funded by Fundação para a Ciência e Tecnologia, respectively with the grants SFRH/BD/38053/2007 and SFRH/BPD/64205/2009. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: fernandoascensao@gmail.com

Introduction

The negative impacts of roads on wildlife have long been recognized [1–5]. Among their many impacts, roads may act as physical barriers to moving animals, thereby reducing landscape connectivity [6,7]. This barrier effect is augmented when wildlife-vehicle collisions (WVC) become significant mortality sources for populations [8,9]. Both WVC and the barrier effect have numerous fitness consequences (e.g., reduced gene flow) that can severely reduce long-term population viability [10,11].

Measures to reduce WVC and mitigate the barrier effect are diverse [12] but wildlife fences combined with crossing structures are gaining more attention by road agencies as they prevent animals from accessing roads while maintaining the connectivity between roadsides [12–14]. However, the choice of mitigation strategy to apply often relies on general patterns, for example road-kill clusters or movement responses [15–17], on the basis that these patterns provide information on the average response of species to roads and traffic. Hence, if individuals vary widely in their responses to roads or mitigation actions and mitigation efforts

are directed toward population-level average responses, these efforts may be only partially effective [18–20].

The life stage and state of an individual can affect its behavioral response to both the road and to the mitigation actions, such as transient individuals avoiding interactions with residents [7,21,22]. For example, squirrel glider (*Petaurus norfolcensis*) movements were re-established across a highway after canopy bridges and glider poles were installed [18,23], yet only half of the individuals known to be present in the vicinity of a canopy bridge used the bridge [18]. The authors suggest that by actively defending their territories, and the passages within them, resident gliders exclude others from accessing those passages [18]. Such behavior could influence monitoring survey results, leading to spurious conclusions on mitigation effectiveness. The importance of thoroughly understanding individual response variability to roads and mitigation is clear.

This knowledge is particularly important for medium-sized carnivores that are especially vulnerable to the negative impacts of roads [8,24,25]. These species typically travel great distances to maintain their territories and occur at low population densities

[26,27]. These traits increase their probability of encountering roads and the significance of each negative interaction. Further, because of their size, these species are often able to trespass through road exclusionary fences, thus being highly exposed to road-kill risk [28–31]. However, it has been shown that these species also regularly use available road passages (e.g., drainage culverts) for safely crossing highways [32–34].

Our goal in this study was to assess whether this apparently contradictory pattern of high road-kill numbers associated with a regular use of road passages is attributable to the variation in behavioral responses toward the highway between individuals. We reanalyzed the tracking data of seven stone martens (*Martes foina*, hereafter referred to as ‘marten’) previously described by [15]. To our knowledge, this is the only available carnivore tracking dataset from a study area that also contains data on road-kill [28] and passage use [33] patterns. As carnivores often occur at low densities, studies investigating their responses to roads often suffer from low sample sizes [35,36], precluding the application of robust analytical methods [37–39]. To overcome small sample size limitations, we employed a novel analytical framework that compares the observed utilization distribution, response turning angles, and highway crossing patterns to results from simulations parameterized with observed data for each individual. We considered these response patterns to describe distinct levels of road impact on marten movements: a greater impact is expected if the utilization distribution across the home range is affected by the highway; an intermediate impact when turning angles are affected by highway proximity; and a more localized impact is expected if crossing patterns are affected by road passage location. Our study design provides a rigorous analytical framework to investigate individual behavior that can be applied across many species and landscapes making it of interest to ecologists, conservation biologists and road planners seeking to understand and mitigate the impacts of roads on carnivore populations.

Materials and Methods

Study area

Martens were tracked in the Mediterranean region of southern Portugal (39°38.154'N, 8°12.128'W), an area dominated by cork-oak woodlands (*Quercus suber*). The study area includes an approximately 10 km section of the four-lane A6 highway and its adjacent surroundings (Fig. 1A). This highway was built in 1995 and has a speed limit of 120 km/h. During the martens activity period in this region (i.e., 20:00 to 08:00 [40]), the A6 receives 169+/-159 vehicles/hour (BRISA S.A., highway enterprise). This highway section has 21 crossing structures available to martens: 13 culverts (1.0–1.5 m in diameter) for water flow, seven underpasses (5 m high, 8 m wide) and one overpass, both for cars and agricultural machinery.

Study species and dataset

The stone marten is a medium-sized carnivore occurring across parts of Asia and Europe [41]. It is often tolerant of human settlements [42], but in our study area it is more commonly associated with cork-oak woodlands [40,43–45]. They are typically solitary and territorial, with home ranges reaching 2 to 3 km² [40]. These martens are particularly sensitive to forest fragmentation [40,43,44,46], and are also highly vulnerable to road-kills, being the second most frequently road-killed carnivore in southern Portugal [28].

We selected seven individuals (two males and five females) that had sufficient data from the dataset used by [15] (Table 1, individual identification herein is that of [15]). These individuals

were tracked between April 2008 and September 2009 during 136 tracking nights (mean 19+/-7 per marten). Each night, one marten was intensively tracked by two observers who attempted to locate the marten every 30 minutes (see [15] for more details). This effort yielded 1425 locations (mean 10+/-5 locations per night per marten) with a mean time between successive locations of 39+/-22 min (Table 1). Pairwise comparisons using Wilcoxon rank sum test with Bonferroni correction revealed a significant higher time interval between relocations in F3 than F2 ($p = 0.008$). However, given the small difference (ca. 5 min.), we do not expect this to preclude the behavior comparison between martens.

Data analysis

Utilization distribution. We estimated marten utilization distributions (UD) with biased random bridges (BRB), a movement-based kernel density estimation method [47,48]. This method improves the spatial resolution of UD estimates by considering activity times between serially correlated relocations rather than simply the spatial density of these relocations as if they were unlinked. The BRB model inserts interpolated locations at regular intervals between each observed location and then uses classical kernel estimation, with a variable smoothing parameter dependent on the time between successive relocations, to estimate the UD [48]. Rather than requiring independence between successive locations, as other space use estimators require [49,50], the BRB model uses the time between successive locations to parameterize the biased random walks between each location [48]. Thus, as the time between successive locations decreases, the width of the bridges (i.e., the size of the area within which the individual may have passed through between successive fixes) decreases, thereby producing a more realistic probability of the animal's true path. Marten space use was estimated within their home range area, which we defined as the 95% isopleth of their UD. BRB were calculated using the ‘BRB’ function within the ‘adehabitatHR’ package (version 0.4.2) [51,52] for R [53].

Movement response angles. We used the nonlinear regression model described by [54] to model the response angles of martens when they approached the highway. The parameters of this model allow us to infer the qualitative response of martens (i.e., attraction, avoidance, or indifference) to the highway proximity. Response angles (A_i) are defined as the difference between the angle of direction to the highway at step S_i and the angle of direction at S_{i+1} , where a step is the estimated path of the animal between successive locations (Fig. S1). These models use the von Mises distribution that is characterized by both the mean angle (μ , angle of maximum probability density) and a concentration parameter (k) that controls the dispersion of the distribution about the mean angle, analogous to the precision of a normal distribution [54]. The distribution is symmetric about the mean angle μ . When $k = 0$, the distribution is uniform on $-\pi$ to π radians. As k increases, the distribution concentrates about the mean angle [54]. We considered two nonlinear models, hereafter referred to as the ‘no-response’ and ‘responsive’ models.

In these models, μ is set constant (i.e., independent of the animal-to-object distance, T_i). In the ‘no-response’ model, k is also independent of the proximity of the object, in this case the highway. Conversely, for the ‘responsive’ model, the concentration parameter is dependent on the animals distance to the highway: the strength of the animal's response is expected to increase with decreasing distance to the highway, so that the animal has a greater tendency to move in the mean response angle μ . As the distance between the animal and the highway increases, the von Mises distribution becomes uniform and the animal's movement




Figure 1. A: Highway A6 in southern Portugal and its crossing structures (squares - culverts, circles - under/over passages), land covers, and marten home range areas (white lines). **B:** Duration (2008–2009) of tracking nights for each marten (each bar is one night) with “LC” indicating loss of contact and “WVC” indicating a confirmed WVC (corpse recovered). Apparent home range overlap of F1 with M1 and M4, and F5 with F7 correspond to distinct periods.
doi:10.1371/journal.pone.0103544.g001

Table 1. Summary data for each marten considered in the present study.

Marten	Number of tracking sessions	Tracking hours	Number of fixes	Mean time between relocations (min)
F1	28	122	202	51 ± 22
F3	28	190	300	44 ± 44
M1	19	137	238	39 ± 22
M4	5	36	64	37 ± 13
F2	19	118	213	39 ± 13
F5	19	124	205	41 ± 16
F7	18	124	203	43 ± 21

Tracked time and time between relocations includes only tracking sessions with at least two successful relocations. Individuals are sorted by whether they crossed the highway (F1, F3, M1, and M4) or not (F2, F5, and F7).
doi:10.1371/journal.pone.0103544.t001

direction becomes more independent of its distance to the highway.

In the ‘responsive’ model the decay of k with the distance to the highway follows an exponential function, governed by two parameters, θ_1 and θ_2 . These two parameters measure the concentration of the von Mises distribution when the distance to the object is zero (θ_1) and the rate of decay of the strength of the animal’s response as it gets farther from the object (θ_2). Statistical inference is likelihood-based, similar to those for generalized linear models to obtain the maximum likelihood estimates [54]. If the highway does not influence marten movement, the ‘no-response’ model will best fit the observed data, whereas the ‘responsive’ model should best fit martens that respond strongly (either attraction or avoidance) to the highway. We excluded locations of inactive martens for this analysis.

Highway crossing patterns. We identified highway crossings as pairs of consecutive marten locations during the same tracking session recorded on opposite sides of the highway. For each marten, we counted the number of crossings and calculated the utilization distribution using these pairs of locations (UD_{cross}) also with the BRB method, similar to the approach used by [55,56].

Null model procedures. We used a null model approach to determine the influence of the highway on marten utilization distribution and highway crossing patterns. Note that the models used to analyze the response angles already incorporate a comparison with a ‘no-response’ model. Null models are pattern-generating simulation models that deliberately exclude a mechanism of interest (for our purposes, the presence of a highway), and by using randomization procedures allow the user to test the importance of that mechanism in observed patterns [57–60]. To build a null model, first an observation is recorded (e.g., number of crossings) from which a set of simulations guided by a set of randomization rules is generated and the simulated response is measured. A large number of iterations (e.g., 1000) are used to generate a frequency histogram of expected response values. The position of the observed response within this null distribution indicates the probability value of the observed pattern, just as in a conventional statistical analysis [58].

Simulated movements were parameterized with the attributes of the observed data (i.e., the number of tracking sessions, locations, step lengths, and utilization distribution boundary), but the simulated agent was naïve to the presence of the highway. For each tracking session, an agent (i.e., simulated marten) started from an observed resting site, chosen randomly, and then moved the same number of steps whose length followed the observed step lengths’ sequence. The agents’ successive location must fall within the home range boundary at a random direction from the previous location. Therefore, simulations follow a constrained random walk which has been successfully used in previous road ecology studies [61–64]. This process is repeated for each tracking session of each marten and each simulated location is saved for further analysis.

For each response considered - utilization distribution, frequency and location of highway crossings - we performed a set of 1000 simulations, per marten. Each set of simulations was used to generate a frequency distribution, from which the confidence intervals of the observed response were estimated. Based on likelihood significance tests, we considered an effect of the highway if the observed parameter fell outside the 5–95% percentiles of the simulated parameter distributions. The model was built using NetLogo 4.1.3 [65] and is available as Model S1.

Influence of land cover on marten movement. Prior to analysis, we investigated marten habitat selection in the study area using a weighted compositional analysis as described by [66]. We

obtained the land cover information by directly classifying Google Earth images using the ‘OpenLayers’ plugin in QGIS (version 1.8) [67]. Ground observations were used to check for and correct potential mismatches. All patches of forest (cork-oak, 67% of the area), agricultural (crop or fallow, 27%), urban (2%), water bodies and streams (1%) present in our study area were polygonized in GIS. Remaining areas were classified as “other” (3%) (Fig. 1A). These land cover polygons were then rasterized to a 30 m grid using the ‘raster’ package (version 2.0-41) [68] for R [53].

For each marten home range, we calculated the sum of the probability values of all cells of the UD for the land cover classes ‘forest’ and ‘agricultural’, and considered these proportions to be the ‘available’ habitat. The ‘used’ habitat was estimated by calculating the proportion of locations that fell within forest or open per marten. The test was performed using the command ‘compana’ in the R package ‘adehabitatHS’(version 2.15.1) [51] with a randomization test (1000 permutations).

Results

Influence of land cover on marten movement

We found no evidence for marten habitat selection ($\Lambda = 0.72$, $p = 0.18$) and so excluded land cover information from further analyses. This was not unexpected, as stone martens in the region are not forest-specialists [40].

Utilization distribution

The UDs of martens whose home ranges overlapped with the highway (F1, F3, M1 and M4) revealed inconsistent patterns of space use near the highway (Fig. 2, left column), with some areas near the highway being used more than expected (Fig. 2, middle column). Less used areas were generally located near territory boundaries and, except for M1, with no relation to highway proximity (Fig. 2, right column).

Movement response angles

Marten responses to the highway were highly variable between individuals. The response angles of five of the seven martens were best predicted by the ‘responsive’ model, suggesting most martens showed a significant response to the highway (Table 2). For example, the mean response angle (μ) of F1 suggested an attraction toward the highway (μ near zero), although the concentration of the von Mises distribution when the distance to the highway approach zero (θ_1) was low (Table 2). Territories of martens F2, F5 and F7 did not overlap with the highway and consequently their mean response angles were high. However, θ_1 was also variable among them, denoting different avoidance levels to the highway proximity. For example, F2 and F5 had θ_1 values of 2.55 and 10.75, respectively, suggesting that the latter had a stronger response to move away from the highway proximity (Table 2). Marten M4 had a mean angle of 1.61, with a θ_1 relatively high, suggesting a predominantly movement parallel to the highway when in its proximity. Interestingly, results for two martens crossing the highway (F3, M1) suggest that their movement was not influenced by highway proximity. The decay of the response angle (θ_2) was low for all martens whose movement was best explained by the ‘responsive’ model, suggesting a nearly linear relation of θ_1 with distance to the highway (Table 2).

Highway crossing patterns

Marten crossing patterns varied between individuals, suggesting no general pattern in highway crossing frequency or crossing locations (Fig. 3). Marten M1 and marginally F1 crossed the highway less often than expected, while M4 crossed more often




Figure 2. Left column: the utilization distributions (UDs, from biased random bridges) of tracked martens with increasing shading indicating increasing use intensity. Marten home-ranges were computed as the 95% isopleth of the UD. Middle and right columns: black areas suggest areas where martens spent more and less time than expected by chance (95% or 5% of simulations), respectively. The highway is shown in each plot by the dotted line. Images are scaled (among martens). Individuals are sorted by whether they crossed the highway (F1, F3, M1 and M4) or not (F2, F5 and F7).

doi:10.1371/journal.pone.0103544.g002

than expected and F3 crossed as often as expected (Fig. 3). For these four martens, their UD_{cross} values suggested that they tended to cross near highway passages. Although most of their UD_{cross} were within the expected interval from the simulations, the highway segments that were used more often than expected have passages (white arrows in Fig. 4). This was particularly clear for F1 and F3. The exception was M4, for which some crossings apparently occurred in sections without any passages (Fig. 4). Interestingly, both F1 and M1 seemed to avoid crossing the highway where paved roads pass beneath the highway (black

arrows in Fig. 4), suggesting a possible behavioral avoidance of this passage type (Fig. 4).

Discussion

Our individual-based analytical framework improved our understanding of how martens respond to the presence of a highway. All martens demonstrated some level of influence of the highway proximity for each of the behavioral responses we considered. However, their responses were more variable than

Table 2. Maximum likelihood estimates of parameters for the ‘responsive’ model: μ - mean angle (in absolute values, ranging from 0 to $|\pi|$ radians); θ_1 - strength of concentration parameter when marten is at distance zero from the highway; θ_2 - rate (exponential) of decay of the concentration parameter as the animal moves farther the highway.

Marten	μ	θ_1	θ_2	χ^2
F1	0.24	0.33	0.000	8.7 (0.00)
F3	1.16	1.32	0.045	2.6 (0.11)
M1	2.30	0.84	0.013	2.3 (0.13)
M4	1.61	1.09	0.008	5.6 (0.02)
F2	2.96	2.55	0.004	6.3 (0.01)
F5	3.08	10.75	0.004	9.3 (0.00)
F7	2.62	6.66	0.005	15.1 (0.00)

Last column stands for the comparison of movement responses to highway proximity, where ‘no-response’ and ‘responsive’ nonlinear models are compared by likelihood ratio test (degrees of freedom = 1). Between brackets is the p-value for the test. Individuals are sorted by whether they crossed the highway (F1, F3, M1, and M4) or not (F2, F5, and F7).

doi:10.1371/journal.pone.0103544.t002

expected [15], highlighting the complexity of individual behavioral toward these linear structures. Because we found no clear evidence for marten habitat selection, we believe that these behavior patterns were mainly due to individual responses to the presence of the highway and crossing structures therein.

We were able to provided new insights into the apparently contradictory results of previous work held in same study area, where martens were frequently killed on the highway [28] while also regularly using crossing structures [33]. We hypothesize that this apparent contradiction could stem from differences between individuals in their familiarity of the highway and the location of passages. For example, seasonal peaks in road mortalities have been well documented elsewhere and coincide with seasonal behavioral patterns, such as breeding behavior, provisioning young or spring dispersal events [28,69,70]. Presumably, these peaks occur because dispersing individuals or individuals exploring new areas in search of mates may be unaware of the passage locations and naïvely cross over the highway, increasing their mortality risk. The support for our hypothesis is described below.

As previously described by [15], some martens maintained territories that overlapped the highway while others maintained

territories adjacent but not overlapping with the highway. However, our results suggest that martens from the former group spent more time than expected in some areas near the highway, while others apparently had no influence of highway proximity on their space use. Linear infrastructures have been used as home range boundaries by carnivores [7] and the martens may have been hunting within the highway verge where prey densities are high [71], although martens with territories adjacent to the highway apparently did not. Hence, we suspect that this high use of the highway verge is due to searching for and using highway crossing structures. This is supported by as the response angles of two martens seem uninfluenced by the highway, which suggests that the highway represented no deterrent for their movement, and they used crossing locations coincident with existing highway crossing structures. Nevertheless, despite the knowledge that these martens regularly cross roads [15], their crossing rates appear highly variable. Moreover, M4 moved parallel to the highway when in its proximity, being the only individual that crossed the highway more often than expected, at both locations coincident and without crossing structures.




Figure 3. Histograms showing the simulated (i.e., predicted) frequency of marten highway crossings (grey bars) and the observed number of crossings (black dot). Grey areas represent the percentile (5–95%) envelope of reference from the simulated datasets. Dots outside of the percentiles suggest the individual crossed less often (left) or more often (right) than expected.

doi:10.1371/journal.pone.0103544.g003




Figure 4. Per marten, top: the utilization distribution of marten highway crossing locations within 200 m from the highway (UD_{cross}). Bottom: the observed probability of crossing the highway at each road segment (UD_{cross} at highway location; solid line). Grey areas represent the 5–95% percentile envelope of reference from the simulated datasets. White (black) arrows indicate highway segments with higher (lower) use than expected. Points indicate road passage location. For each marten, the highway segment in the upper-half of the figure is projected in the X axis from the bottom picture.
doi:10.1371/journal.pone.0103544.g004

Overall, although the sample size of our dataset is less than ideal, we show that martens can exhibit a variety of responses to the highway, especially in their propensity to cross the highway and to use crossing structures to do so. We assume that martens F1, F2 and F3 were residents with well-established territories since they were tracked for long periods, having stable home range areas [15], and F1 and F3 were apparently aware of passage locations for crossings. The movement of these two martens was not hindered by the presence of the highway probably because they knew where to access suitable crossing structures. Although we did not monitor the existing road passages, *ad-hoc* observations confirmed that F3 regularly used the structure with higher probability of use (author's *pers. obs.*, passage below the white arrow in Fig. 4).

We also believe that M4 was dispersing through the region as he was not detected before being captured, despite the continuous and intense trapping effort [15]. This marten was trapped in November when martens typically disperse [72,73] and was road-killed shortly after arriving to the study area. Assuming this marten was not a resident he would have been naïve to the location of

passages or could have been prevented from accessing them. This would explain that, unlike other martens, some crossings of M4 occurred in sections without passages. Our assumption that non-residents are unaware of the passage locations for safe crossings is supported by previous work suggesting that individuals require time to adapt to existing crossing structures [74–77]. Thus, our findings suggest that the use of passages seems to be governed not only by road and environmental attributes [78], but also by individual preferences and familiarity with the landscape.

To effectively mitigate the negative effects of roads at the population level we must understand the processes that affect the movements of individuals and the variability between individual responses to roads and existing mitigation [16]. For example, marten use of crossing structures, particularly culverts, is well documented [32–34,79,80]. However, our results suggest individual preferences for specific crossing structures: F3 crossed the highway at least 30 times during our tracking sessions, but apparently did so through a single passage despite at least two similar structures being within her home range (Fig. 4). Additionally, F1 and M1 both used and avoided the same passages (white

and black arrows, respectively, in Fig. 4). These differences may reflect individual preferences for passage characteristics, locations, or both.

An important research question remains: how many individuals use the passages? If only a few individuals regularly use the same passage, as our results suggest, then the effectiveness of these structures could be overestimated [18]. This topic is of major importance [12], yet poorly understood [but see 18,81]. This information, together with a deeper understanding of animal behavior near roads, would provide a spatio-temporal bridge between the individual and its population [82,83]. Mitigation strategies that ignore this information may be insufficient. Given the amount of time these martens spend near the highway, the variety of responses they demonstrated toward the highway, and their apparent passage-type preferences, we believe mitigation strategies would be more effective by optimizing the number of and spacing between passages [84], and by directing new individuals toward existing passages via exclusionary fencing with a sufficiently small mesh size as previously suggested [85,86]. Such

mitigation measures are necessary to reduce the mortality risks to both resident and importantly to dispersing individuals.

Supporting Information

Figure S1 Diagram illustrating the response angle in relation to the highway location (grey line). The animal moves from S_i to S_{i+1} . The animal-to-highway angle in radians is C_i . The move angle is B_i and the response angle is A_i . (DOCX)

Model S1 NetLogo model built to simulate marten movement in highway vicinity. (NLOGO)

Author Contributions

Conceived and designed the experiments: FA. Performed the experiments: FA. Analyzed the data: FA. Contributed reagents/materials/analysis tools: FA JT. Wrote the paper: FA CG SLP JT AC MSR. Responsible for the dataset analyzed: CG.

References

- Forman RTT, Sperling D, Bissonette J, Clevenger A, Cutshall C, et al. (2003) Road ecology: science and solutions. Washington, DC: Island Press.
- Beckmann JP, Clevenger AP, Huijser M, Hilty JA (2010) Safe passages: highways, wildlife, and habitat connectivity. Island Press.
- Trombulak SC, Frissell CA (2000) Review of ecological effects of roads on terrestrial and aquatic communities. *Conservation Biology* 14: 18–30.
- Forman RTT, Alexander LE (1998) Roads and their major ecological effects. *Annual Review of Ecology and Systematics* 29: 207–231.
- Stoner D (1925) The Toll of the Automobile. *Science* 61: 56–57.
- Jackson ND, Fahrig L (2011) Relative effects of road mortality and decreased connectivity on population genetic diversity. *Biological Conservation* 144: 3143–3148.
- Riley SPD, Pollinger JP, Sauvajot RM, York EC, Bromley C, et al. (2006) A southern California freeway is a physical and social barrier to gene flow in carnivores. *Molecular Ecology* 15: 1733–1741.
- Benítez-López A, Alkemade R, Verweij PA (2010) The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biological Conservation* 143: 1307–1316.
- Fahrig L, Rytwinski T (2009) Effects of Roads on Animal Abundance: an Empirical Review and Synthesis. *Ecology and Society* 14: 21. URL: <http://www.ecologyandsociety.org/vol14/iss21/art21/>.
- Eigenbrod F, Hecnar SJ, Fahrig L (2008) Accessible habitat: an improved measure of the effects of habitat loss and roads on wildlife populations. *Landscape Ecology* 23: 159–168.
- Holderegger R, Di Giulio M (2010) The genetic effects of roads: A review of empirical evidence. *Basic and Applied Ecology* 11: 522–531.
- van der Grift EA, van der Ree R, Fahrig L, Findlay S, Houlihan J, et al. (2013) Evaluating the effectiveness of road mitigation measures. *Biodiversity and Conservation* 22: 425–448.
- Iuell B, Bekker GJ, Cuperus R, Dufek J, Fry G, et al. (2003) COST 341 - Wildlife and Traffic: A European Handbook for Identifying Conflicts and Designing Solutions. URL: <https://www.milieuinfo.be/productie/beheerplone/nietaem/ene/cost-341/COST%20341-handbook.pdf> (assessed at 28-10-2012).
- Glista DJ, DeVault TL, DeWoody JA (2009) A review of mitigation measures for reducing wildlife mortality on roadways. *Landscape Urban Plan* 91: 1–7.
- Grilo C, Sousa J, Ascensão F, Matos H, Leitão I, et al. (2012) Individual Spatial Responses towards Roads: Implications for Mortality Risk. *PLoS ONE* 7: e43811.
- Klar N, Fernandez N, Kramer-Schadt S, Herrmann M, Trinzen M, et al. (2008) Habitat selection models for European wildcat conservation. *Biological Conservation* 141: 308–319.
- Gunson KE, Mountrakis G, Quackenbush IJ (2011) Spatial wildlife-vehicle collision models: A review of current work and its application to transportation mitigation projects. *Journal of Environmental Management* 92: 1074–1082.
- Soanes K, Lobo MC, Vesik PA, McCarthy MA, Moore JL, et al. (2013) Movement re-established but not restored: Inferring the effectiveness of road-crossing mitigation for a gliding mammal by monitoring use. *Biological Conservation* 159: 434–441.
- Corlatti L, Hacklauer K, Frey Roos F (2009) Ability of wildlife overpasses to provide connectivity and prevent genetic isolation. *Conservation Biology* 23: 548–556.
- Simmons JM, Sunnucks P, Taylor AC, van der Ree R (2010) Beyond road-kill, radiotracking, recapture and FSTs a review of some genetic methods to improve understanding of the influence of roads on wildlife. *Ecology & Society* 15: 9–24.
- Laundre JW, Hernandez L, Altendorf KB (2001) Wolves, elk, and bison: reestablishing the “landscape of fear” in Yellowstone National Park, USA. *Can J Zool* 79: 1401–1409.
- Valeix M, Loveridge AJ, Chamaille-James S, Davidson Z, Murindagomo F, et al. (2009) Behavioral adjustments of African herbivores to predation risk by lions: Spatiotemporal variations influence habitat use. *Ecology* 90: 23–30.
- van der Ree R, Cesarini S, Sunnucks P, Moore JL, Taylor A (2010) Large Gaps in Canopy Reduce Road Crossing by a Gliding Mammal. *Ecology and Society* 15.
- Rytwinski T, Fahrig L (2012) Do species life history traits explain population responses to roads? A meta-analysis. *Biological Conservation* 147: 87–98.
- Rytwinski T, Fahrig L (2013) Why are some animal populations unaffected or positively affected by roads? *Oecologia* 173: 1143–1156.
- Crooks KR (2002) Relative Sensitivities of Mammalian Carnivores to Habitat Fragmentation. *Conservation Biology* 16: 488–502.
- Noss RF, Quigley HB, Hornocker MG, Merrill T, Paquet PC (1996) Conservation biology and carnivore conservation in the Rocky Mountains. *Conservation Biology* 10: 949–963.
- Grilo C, Bissonette JA, Santos-Reis M (2009) Spatial-temporal patterns in Mediterranean carnivore road casualties: Consequences for mitigation. *Biological Conservation* 142: 301–313.
- Clarke GP, White PCL, Harris S (1998) Effects of roads on badger *Meles meles* populations in south-west England. *Biological Conservation* 86: 117–124.
- Hauer S, Ansohn H, Zinke O (2002) Mortality patterns of otters (*Lutra lutra*) from eastern Germany. *Journal of Zoology* 256: 361–368.
- Philcox CK, Grogan AL, Macdonald DW (1999) Patterns of otter *Lutra lutra* road mortality in Britain. *Journal of Applied Ecology* 36: 748–761.
- Ascensão F, Mira A (2007) Factors affecting culvert use by vertebrates along two stretches of road in southern Portugal. *Ecological Research* 22: 57–66.
- Grilo C, Bissonette JA, Santos-Reis M (2008) Response of carnivores to existing highway culverts and underpasses: implications for road planning and mitigation. *Biodiversity and Conservation* 17: 1685–1699.
- Serronha AM, Mateus AR, Eaton F, Santos-Reis M, Grilo C (2012) Towards effective culvert design: monitoring seasonal use and behavior by Mediterranean mesocarnivores. *Environmental Monitoring and Assessment* (online first 2012/12/05).
- LaPoint S, Gallery P, Wikelski M, Kays R (2013) Animal behavior, cost-based corridor models, and real corridors. *Landscape Ecology* 28: 1615–1630.
- Herr J, Schley L, Roper TJ (2009) Socio-spatial organization of urban stone martens. *Journal of Zoology* 277: 54–62.
- Roedenbeck IA, Fahrig L, Findlay CS, Houlihan JE, Jaeger JAG, et al. (2007) The Rauschholzhausen agenda for road ecology. *Ecology and Society* 12: 11. URL: <http://www.ecologyandsociety.org/vol12/iss11/art11/>.
- van der Ree R, Jaeger JAG, van der Grift EA, Clevenger AP (2011) Effects of Roads and Traffic on Wildlife Populations and Landscape Function: Road Ecology is Moving toward Larger Scales. *Ecology & Society* 16.
- Lesbarreres D, Fahrig L (2012) Measures to reduce population fragmentation by roads: what has worked and how do we know? *Trends Ecol Evol* 27: 374–380.
- Santos-Reis M, Santos M, Lourenço S, Marques J, Pereira I, et al. (2004) Relationships between Stone Martens, Genets and Cork Oak Woodlands in Portugal. In: D. J. Harrison, A. K. Fuller and G. Proulx, editors. *Martens and Fishers (Martes) in Human-Altered Environments An International Perspective*. Massachusetts: Springer-Verlag, pp. 147–172.
- Proulx G, Aubry K, Birks J, Buskirk S, Fortin C, et al. (2005) World distribution and status of the genus *Martes* in 2000. In: D. J. Harrison, A. K. Fuller and

- G. Proulx, editors. Martens and Fishers (Martes) in Human-Altered Environments. New York: Springer. pp. 21–76.
42. Herr J, Schley L, Roper TJ (2009) Stone martens (*Martes foina*) and cars: investigation of a common human–wildlife conflict. *European Journal of Wildlife Research* 55: 471–477.
 43. Mortelliti A, Boitani L (2008) Interaction of food resources and landscape structure in determining the probability of patch use by carnivores in fragmented landscapes. *Landscape Ecology* 23: 285–298.
 44. Grilo C, Ascensão F, Santos-Reis M, Bissonette JA (2011) Do well-connected landscapes promote road-related mortality? *European Journal of Wildlife Research* 57: 707–716.
 45. Santos MJ, Santos-Reis M (2010) Stone marten (*Martes foina*) habitat in a Mediterranean ecosystem: effects of scale, sex, and interspecific interactions. *European Journal of Wildlife Research* 56: 275–286.
 46. Virgós E, Tellería JL, Santos T (2002) A comparison on the response to forest fragmentation by medium-sized Iberian carnivores in central Spain. *Biodiversity and Conservation* 11: 1063–1079.
 47. Benhamou S, Cornéris D (2010) Incorporating movement behavior and barriers to improve kernel home range space use estimates. *The Journal of Wildlife Management* 74: 1353–1360.
 48. Benhamou S (2011) Dynamic Approach to Space and Habitat Use Based on Biased Random Bridges. *PLoS ONE* 6 e14592.
 49. Swihart RK, Slade NA (1985) Influence of Sampling Interval on Estimates of Home-Range Size. *Journal of Wildlife Management* 49: 1019–1025.
 50. Worton BJ (1989) Kernel methods for estimating the utilization distribution in home-range studies. *Ecology* 70: 164–168.
 51. Calenge C (2006) The package “adehabitat” for the R software: A tool for the analysis of space and habitat use by animals. *Ecological Modelling* 197: 516–519.
 52. Calenge C (2011) Home Range Estimation in R: the adehabitatHR Package. URL: <http://cran.r-project.org/web/packages/adehabitatHR/vignettes/adehabitatHR.pdf>.
 53. R-Core-Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
 54. Tracey J, Zhu J, Crooks K (2005) A set of nonlinear regression models for animal movement in response to a single landscape feature. *Journal of Agricultural, Biological, and Environmental Statistics* 10: 1–18.
 55. Horne JS, Garton EO, Krone SM, Lewis JS (2007) Analyzing animal movements using Brownian bridges. *Ecology* 88: 2354–2363.
 56. Lewis JS, Rachlow JL, Horne JS, Garton EO, Wakkinen WL, et al. (2011) Identifying habitat characteristics to predict highway crossing areas for black bears within a human-modified landscape. *Landscape Urban Plan* 101: 99–107.
 57. Gotelli NJ (2001) Research frontiers in null model analysis. *Global Ecology and Biogeography* 10: 337–343.
 58. Gotelli NJ, Graves GR (1996) Null models in ecology. Washington: Smithsonian Institution Press.
 59. Martin J, Calenge C, Quenette P-Y, Allainé D (2008) Importance of movement constraints in habitat selection studies. *Ecological Modelling* 213: 257–262.
 60. Richard E, Calenge C, Saïd S, Hamann JL, Gaillard JM (2012) Studying spatial interactions between sympatric populations of large herbivores: a null model approach. *Ecography* 36: 157–165.
 61. Beyer HL, Ung R, Murray DL, Fortin MJ (2013) Functional responses, seasonal variation and thresholds in behavioural responses of moose to road density. *Journal of Applied Ecology* 50: 286–294.
 62. Rondinini C, Boitani L (2002) Habitat use by beech martens in a fragmented landscape. *Ecography* 25: 257–264.
 63. Shepard DB, Kuhns AR, Dreslik MJ, Phillips CA (2008) Roads as barriers to animal movement in fragmented landscapes. *Animal Conservation* 11: 288–296.
 64. Whittington J, St Clair CC, Mercer G (2004) Path tortuosity and the permeability of roads and trails to wolf movement. *Ecology and Society* 9: 4. URL: <http://www.ecologyandsociety.org/vol9/iss1/art4/>.
 65. Wilensky U (1999) NetLogo: Center for connected learning and computer-based modeling. Evanston, IL: Northwestern University.
 66. Millsbaugh JJ, Nielson RM, McDonald L, Marzluff JM, Gitzen RA, et al. (2006) Analysis of resource selection using utilization distributions. *Journal of Wildlife Management* 70: 334–395.
 67. Quantum-GIS-Development-Team (2011) Quantum GIS Geographic Information System. Open Source Geospatial Foundation. Available: <http://qgis.osgeo.org>.
 68. Hijmans RJ, van Etten J (2012) raster: Geographic analysis and modeling with raster data. R package version 2.0-12. <http://CRAN.R-project.org/package=raster>.
 69. Morelle K, Lechaire F, Lejeune P (2013) Spatio-temporal patterns of wildlife-vehicle collisions in a region with a high-density road network. *Nature Conservation* 5: 53–73.
 70. Steiner W, Leisch F, Hackländer K (2014) A review on the temporal pattern of deer–vehicle accidents: Impact of seasonal, diurnal and lunar effects in cervids. *Accident Analysis & Prevention* 66: 168–181.
 71. Ascensão F, Clevenger AP, Grilo C, Filipe J, Santos-Reis M (2012) Highway verges as habitat providers for small mammals in agrosilvopastoral environments. *Biodiversity and Conservation* 21: 3681–3697.
 72. Mangas J (2009) Garduña – Martes foina. In: A. Salvador and J. Cassinello, editors. *Enciclopedia Virtual de los Vertebrados Españoles*. Museo Nacional de Ciencias Naturales, Madrid. URL: <http://www.vertebradosibericos.org/>.
 73. Blanco JC, González JL (1992) *Livro Rojo de Los Vertebrados de España*. Madrid: Ministerio de la Agricultura, Pesca y Alimentación, ICONA.
 74. Clevenger AP, Ford AT, Sawaya MA (2009) Banff wildlife crossings project: Integrating science and education in restoring population connectivity across transportation corridors. Final report to Parks Canada Agency, Radium Hot Springs, British Columbia, Canada. pp. 165.
 75. Dodd NL, Gagnon JW, Manzo AL, Schweinsburg RE (2007) Video surveillance to assess highway underpass use by elk in Arizona. *Journal of Wildlife Management* 71: 637–645.
 76. Gagnon JW, Dodd NL, Ogren KS, Schweinsburg RE (2011) Factors Associated With Use of Wildlife Underpasses and Importance of Long-Term Monitoring. *Journal of Wildlife Management* 75: 1477–1487.
 77. Clevenger AP, Walther N (2005) Performance indices to identify attributes of highway crossing structures facilitating movement of large mammals. *Biological Conservation* 121: 453–464.
 78. Clevenger AP, Chruszcz B, Gunson K (2001) Drainage culverts as habitat linkages and factors affecting passage by mammals. *Journal of Applied Ecology* 38: 1340–1349.
 79. Mateus ARA, Grilo C, Santos-Reis M (2011) Surveying drainage culvert use by carnivores: sampling design and cost-benefit analyses of track-pads vs. video-surveillance methods. *Environmental monitoring and assessment* 181: 101–109.
 80. Rodríguez A, Crema G, Delibes M (1996) Use of non-wildlife passages across a high speed railway by terrestrial vertebrates. *Journal of Applied Ecology* 33: 1527–1540.
 81. Sawaya MA, Stetz JB, Clevenger AP, Gibeau ML, Kalinowski ST (2012) Estimating Grizzly and Black Bear Population Abundance and Trend in Banff National Park Using Noninvasive Genetic Sampling. *PLoS ONE* 7.
 82. Patterson TA, Thomas L, Wilcox C, Ovaskainen O, Matthiopoulos J (2008) State-space models of individual animal movement. *Trends in Ecology & Evolution* 23: 87–94.
 83. Schick RS, Loarie SR, Colchero F, Best BD, Boustany A, et al. (2008) Understanding movement data and movement processes: current and emerging directions. *Ecology Letters* 11: 1338–1350.
 84. Bissonette JA, Adair W (2008) Restoring habitat permeability to roaded landscapes with isometrically-scaled wildlife crossings. *Biological Conservation* 141: 482–488.
 85. Ascensão F, Clevenger A, Santos-Reis M, Urbano P, Jackson N (2013) Wildlife–vehicle collision mitigation: Is partial fencing the answer? An agent-based model approach. *Ecological Modelling* 257: 36–43.
 86. Klar N, Herrmann M, Kramer-Schadt S (2009) Effects and Mitigation of Road Impacts on Individual Movement Behavior of Wildcats. *Journal of Wildlife Management* 73: 631–638.